arXiv:math/0103015v3 [math.MG] 22 Aug 2001

SINGULAR SETS AND PARAMETERS OF GENERALIZED TRIANGLE ORBIFOLDS

MICHAEL BELOLIPETSKY

Abstract. We study groups generated by three half-turns in Lobachevsky space and their factor-orbifolds. This generalized triangle groups are directly connected with the arbitrary 2-generator Kleinian groups. Our main result is a complete description of the singular sets of the generalized triangle orbifolds. We also give a method to obtain the parameters defining a generalized triangle group from the structure of the singular set of its factor-orbifold and illustrate it by examples.

1. Introduction

After the famous lectures of W. Thurston [28] the study of 3-dimensional manifolds was focused on geometric and especially on hyperbolic manifolds and orbifolds. An orientable hyperbolic 3-orbifold can be obtained as a factor-space of the Lobachevsky space $\mathcal{H}^3$ by the action of a discrete subgroup of the group of isometries $Isom^+\mathcal{H}^3 = \mathrm{PSL}(2,\mathbb{C})$. So the hyperbolic 3-orbifolds are connected with the discrete subgroups of $\mathrm{PSL}(2,\mathbb{C})$ known also as *Kleinian groups*. Since a general subgroup of $\mathrm{PSL}(2,\mathbb{C})$ is discrete if and only if each of its 2-generator subgroups is discrete (see e.g. [4]), the class of the 2-generator Kleinian groups takes on special significance. This groups and in particular arithmetic 2-generator Kleinian groups were studied by F. Gehring, C. Maclachlan, G. Martin, J. Montesinos, A. Reid and others (see [5], [7], [12], [14], [15], [19], [21], [25] and the references therein). It appears, that the 3-orbifolds corresponding to the 2-generator Kleinian groups may have a very complicated structure. In a special case of the groups with real parameters the orbifolds were studied by J. Gilman [13], E. Klimenko [22],[23]; and have been recently completely classified by E. Klimenko and N. Kopteva (see [24]).

In this article we consider the groups of hyperbolic isometries generated by three half-turns in $\mathcal{H}^3$. If the axes of the half-turns pairwise intersect then the group is isomorphic to a Fuchsian triangle group, so the groups generated by three half-turns can be considered as a generalization of the Fuchsian triangle groups. The generalized triangle groups are directly connected with the arbitrary 2-generator Kleinian groups (see Section 5) but have much more trackable geometric structure. Thus, since all three generators of the groups have fixed points (axes) in $\mathcal{H}^3$, the fundamental groups of underlying spaces of the corresponding 3-orbifolds are always trivial, so the underlying space of such an orbifold is a 3-sphere $\mathcal{S}^3$ or a counterexample to the Poincare conjecture.

Further we will always suppose that the underlying space is $\mathcal{S}^3$. In Section 2 we describe singular structures of common generalized triangle orbifolds, which are naturally fall into eight types (Theorem 2.4).

The set of the generalized triangle groups can be parameterized by three complex numbers that correspond to the complex distances [9] between the axes of the generators. In Section 3 we use W. Fenchel's technics [9] to study this parameterization and its connection with the matrix representation of the group in $\mathrm{SL}(2, \mathbb{C})$. Section 4 shows how to find the parameters of the group for the eight types of singular structures from Theorem 2.4. The connection between the generalized triangle groups and the 2-generator Kleinian groups and their usual parameterization is considered in Section 5.

Section 6 presents examples of common generalized triangle orbifolds with different types of singularities and in Section 7 we consider an example for the exceptional case.

Acknowledgement. I would like to thank Professor Alexander D. Mednykh for interesting me in this problem and numerous helpful discussions.

2. Structure of the singular sets

The underlying space of a generalized triangle group is 3-sphere, so the singular sets are knotted graphs in $\mathcal{S}^3$ with the orbifolds fundamental groups generated by three involutions. While depicting the singular sets we follow a common notation writing indexes of singularity near corresponding components and omitting indexes 2. We will use Wirtinger presentation of the fundamental group of a knotted graph [10] adopted to the group of orbifold [16], so the elementary arcs of the graph will correspond to the words in the group.

Definition 2.1. Common generalized triangle group *is a Kleinian group generated by three halftuns such that the fundamental group of the complement of its singular set in* $\mathcal{S}^3$ *has also rank* 3.

This definition allows us in some cases to consider groups of the complements of the singular graphs instead of the orbifold groups. This saves from certain difficulties, which may arise because the considered groups have many involutions.

Lemma 2.1. *The class of the common generalized triangle groups is invariant under Reidemeister moves for graphs.*

Proof. This lemma is obvious because the Reidemeister moves do not change the fundamental group of the complement of the singular graph. □

Now suppose we have a knotted graph such that some three arcs of the graph correspond to the three elements generating the fundamental group of the complement of the graph in $\mathcal{S}^3$. Then corresponding three involutions generate our generalized triangle group. The first thing we do is we draw a circle that contains a plane projection of the singular graph and using the Reidemeister moves draw the generating arcs outside the circle:



Figure 1. The singular set (except the generating arcs) of generalized triangle group $< a, b, c >$ is in disc D.

Here we have to note that Lemma 2.1 is not enough to carry on this procedure because it does not say anything about the generators of the group, and what we want is to obtain the group generated by the three elements that correspond to the selected singular arcs. However, it is a matter of a simple case-by-case verification to see that the generators can be "preserved" and we leave it as an exercise.

Now let us see what singular structures are possible inside D. The only restrictions we have are that the group should be generated by a, b, c and that the corresponding factor-orbifold is hyperbolic.

Lemma 2.2. *Non trivial knots on singular arcs inside D are impossible.*



Figure 2. An illustration for the proof of Lemma 2.2.

Proof. Suppose there is a non-trivial knot on a singular arc inside D. Let X be the complement to the singular set in S^3. Consider decomposition $X = X_1 \cup X_2$ such that X_2 contains the knot and ∂X_2 has precisely 2 punctures (see Figure 2). Here X_2 could be obtained from the tubular neighborhood of the singular set, so the decomposition is always possible.

We now apply Van Kampen theorem to $X = X_1 \cup X_2$. For $P \in X_1 \cap X_2$:

$$\pi_1(X_1, P) =< a, b, c \mid \cdots >,$$

$$\pi_1(X_2, P) =< w_1', w_2', x, \cdots \mid \cdots >$$

and x can not be obtained as a word of w_1', w_2' since the knot is non-trivial.

By Van Kampen theorem

$$\pi_1(X, P) =< a, b, c, w_1', w_2', x, \cdots \mid w_1 w_1', w_2 w_2' \cdots >=< a, b, c, x, \cdots \mid \cdots > .$$

Here again x can not be obtained as a word of the other generators. This means that the rank of $\pi_1(X)$ is greater then 3, which contradicts the definition of the common generalized triangle group. □

Since arcs of the graph can not be knotted with themselves, the singular graph has a stratiform structure shown on Figure 3 with all the vertices inside D_1 (we use the common notation for integer tangles with l_i, m_i, $n_i \in \mathbb{Z}$ for $i = 1, \ldots, k$).



FIGURE 3. Structure of the singular set given by signature $S(l_1, m_1, n_1, l_2, m_2, n_2, \ldots, l_k, m_k, n_k)$.

Lemma 2.3. *The singular graph in D_1 has no cycles.*

Proof. Let X be the common generalized triangle orbifold, and $\tilde{X}$ - the 3-manifold obtained as the complement to the singular set of X in S^3. To prove the lemma it is sufficient to consider the first homology group $H_1(\tilde{X})$. The canonical projection $\tilde{p} : \pi_1(\tilde{X}) \to \pi_1(\tilde{X})^{\mathrm{ab}} \cong H_1(\tilde{X})$ maps non of the generators of $\pi_1(\tilde{X})$ to the homology cycle that corresponds to the singular cycle in D_1. Since $H_1(\tilde{X}) \cong \mathbb{Z}^r$ with no any elements of finite order, this immediately follows that $rank(H_1(\tilde{X})) > rank(\tilde{p}(\pi_1(\tilde{X})))$, which is impossible.

Note, that this argument does not work for the orbifold fundamental and homology groups because the homology groups are generated by the elements of the finite order (See example in Section 7). ☐

Lemma 2.3 makes it possible to push all remaining tangles out of D_1 by drawing vertices through the tangles as it shown on Figure 4.



FIGURE 4. Drawing a vertex through a tangle.

So for the graph in D_1 we have:

- it has no cycles;
- it has no tangles;
- the group of its complement is generated by (six) arcs coming into D_1 from the outside.

Together with the assumption that the orbifold is hyperbolic this gives all we need to classify singular sets of the common generalized triangle groups. In the following theorem we summarize the results:

Theorem 2.4. *The structure of the singular set of a common generalized triangle orbifold is determined by the signature* $S(l_1, m_1, n_1, l_2, m_2, n_2, \ldots, l_k, m_k, n_k)$ *(Figure 3) together with* D_1 *corresponding to one of the cases A - H (Figure 5).*



FIGURE 5. Singular structures inside D_1 (here $t_i \in \mathbb{N} \cup \{\infty\}$).

Orbifolds obtained from A, B, C (with $1/t_1+1/t_2+1/t_3 < 1$*), D (with* $1/t_1+1/t_2 < 1/2$ *and* $1/t_2 + 1/t_3 < 1/2$*) are compact; while orbifolds obtained from C (with*

$1/t_1+1/t_2+1/t_3=1$), D (with $1/t_1+1/t_2=1/2$ or $1/t_2+1/t_3=1/2$), E, F, G and H are non compact.

Remark 1. We have dropped one more combinatorially possible case that can be obtained from G deleting the arc, which joins the vertices, because corresponding groups are always generated by only two half-turns.

Remark 2. Some relatively simple singular structures from Theorem 2.4 correspond to non-hyperbolic orbifolds (see Examples 6C, 6D). All such geometric orbifolds can be found in [8] and we do not specially focus on non-hyperbolic case.

Remark 3. A generalized triangle orbifold is uniquely determined by the signature and D_1 as it was shown above, however different signatures can correspond to the same orbifolds.

3. Matrix representation and parameters

In order to obtain a matrix representation for a group generated by three half-turns one has to fix the axes of the generators. We do this by means of complex distances between the hyperbolic lines ([9], p. 67-70). We define the *complex distance* between two oriented hyperbolic lines as follows: its real part equals the length of the common perpendicular to the lines and imaginary part is given by the angle between the lines, taken from the first to the second line respecting orientations.

We call three complex distances between the axes of the generators by the *parameters* of the group $\Gamma=\langle a,b,c\rangle$. Let us remark that starting from this place by group $\langle a,b,c\rangle$ we mean the group with the given generators taken in a fixed order, so, in fact, we work with a *marked groups* generated by a,b,c.

$$par(\langle a,b,c\rangle)=(\mu(a,b),\ \mu(a,c),\ \mu(c,b))=(\mu_0,\ \mu_1,\ \mu_2).$$

(Here $\mu(\dots,\dots)=x+iy$, $x\in\mathbb{R}^+$, $y\in[0;2\pi)$ — the complex distance between the oriented axes of the half-turns.)

Remark. Since the directions of the half-turns are undefined the parameters are defined up to orientation of the axes. We can canonically link the orientation of a hyperbolic line with the matrix of the half-turn in $\mathrm{SL}(2,\mathbb{C})$ (see [9], p. 63). This means that fixing the directions of the axes of generators equals choosing the inverse image after canonical projection $p:\mathrm{SL}(2,\mathbb{C})\to\mathrm{PSL}(2,\mathbb{C})$ of our $\mathrm{PSL}(2,\mathbb{C})$-subgroup in $\mathrm{SL}(2,\mathbb{C})$. Following common agreement we usually call this inverse image by representation of our group in $\mathrm{SL}(2,\mathbb{C})$.

Lemma 3.1. *Complex vector $par(\langle a,b,c\rangle)$ determines group $\langle a,b,c\rangle$ uniquely up to conjugation in $\mathrm{PSL}(2,\mathbb{C})$, i.e. if $par(\langle a,b,c\rangle)=par(\langle a',b',c'\rangle)$ then there exists $h\in\mathrm{PSL}(2,\mathbb{C})$ such that $a'=hah^{-1}$, $b'=hbh^{-1}$, $c'=hch^{-1}$.*

Proof. Consider the axes of the half-turns a,b,c in $\mathcal{H}^3$. We endow the axes by orientations and join them in couples by common perpendiculars. The result is a hyperbolic right-angled hexagon (Figure 6).



FIGURE 6. Axes of the half-turns a, b, c together with their common perpendiculars give a hyperbolic right-angled hexagon.

Vector $par(\langle a,b,c\rangle)$ represents three complex lengths of the sides of our hexagon which are common perpendiculars of the given axes. By theorem from [9] (p. 94) three pairwise non-adjacent sides define right-angled hexagon uniquely up to simultaneous change of orientations of the other three sides. It means that any other hexagon defined by another group with the same parameters can be translated to the given one by a hyperbolic isometry. This isometry obviously defines the required conjugation $h \in \mathrm{PSL}(2,\mathbb{C})$.

It may seem that we have missed the possibility of changing of the orientations of the axes but we have already remarked that the orientations effect only representation of the group in $\mathrm{SL}(2,\mathbb{C})$ but not isometries themselves. So in this situation we can change the orientations as it is needed. ☐

Next thing to do is to find out some convenient representation for $\langle a,b,c\rangle$ in $\mathrm{SL}(2,\mathbb{C})$ and describe it in terms of $par(\langle a,b,c\rangle)$. We denote the matrices corresponding to the half-turns in a, b, c by capitals A, B, C.

Half-turn in oriented hyperbolic line m is represented by a matrix $M \in \mathrm{SL}(2,\mathbb{C})$ of the form $\left(\begin{array}{cc} m_{11} & m_{12} \\ m_{21} & -m_{11} \end{array}\right)$, which is characterized by $M^2 = -I$ or equivalently $\mathrm{tr(M)} = 0$. This matrix M is called *the normalized line-matrix* of line m. By extending transformation M to the absolute $\overline{\mathbb{C}} = \partial\mathcal{H}^3$ it is easy to find the set of its fixed points in $\overline{\mathbb{C}}$ denoted by $fix(M)$. Geometrically the fixed points $\{z_1,\ z_2\} = fix(M)$ are the ends of the arc orthogonal to $\mathbb{C}$ which presents line m in the upper-space model of $\mathcal{H}^3$.

After a suitable conjugation of the group $\langle A,B,C\rangle$ in $\mathrm{SL}(2,\mathbb{C})$ one can suppose that $fix(A) = \{1/\beta; -1/\beta\}$, $fix(B) = \{\beta; -\beta\}$ ($\beta \in \mathbb{C}$ and $|\beta| \geq 1$).

We have:

$$A = \begin{pmatrix} 0 & i/\beta \\ i\beta & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & i\beta \\ i/\beta & 0 \end{pmatrix}, \quad C = \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & -c_{11} \end{pmatrix},$$

where $-c_{11}^2 + c_{12}c_{21} = 1$ since $C \in \mathrm{SL}(2,\mathbb{C})$.

To find β and c_{ij} in terms of $par(\langle a,b,c\rangle)$ we use the following important formula, which allows to find the complex distance between hyperbolic lines m_1, m_2 from the matrices M_1, M_2 of the lines ([9], p. 68):

$$cosh(\mu(m_1,m_2)) = -\frac{1}{2}\mathrm{tr}\,(\mathrm{M_1M_2}).$$

Denote $\rho_k = -2\,cosh\mu_k$ $(k = 0,\ 1,\ 2)$. Then

$$\rho_0 = \mathrm{tr(AB)} = \mathrm{tr}\begin{pmatrix} -1/\beta^2 & 0 \\ 0 & -\beta^2 \end{pmatrix};$$

$$\rho_1 = \mathrm{tr(AC)} = \mathrm{tr}\begin{pmatrix} ic_{21}/\beta & -ic_{11}/\beta \\ ic_{11}\beta & ic_{12}\beta \end{pmatrix};$$

$$\rho_2 = \mathrm{tr(BC)} = \mathrm{tr}\begin{pmatrix} ic_{21}\beta & -ic_{11}\beta \\ ic_{11}/\beta & ic_{12}/\beta \end{pmatrix}.$$

We obtain system of equations on β and c_{ij}:

$$\begin{cases} -1/\beta^2 - \beta^2 = \rho_0; \\ ic_{21}/\beta + ic_{12}\beta = \rho_1; \\ ic_{21}\beta + ic_{12}/\beta = \rho_2; \\ c_{11} = i\sqrt{c_{12}c_{21}+1}. \end{cases}$$

Solving this equations we find out all parameters in our representation. Let us remark that this system has not unique solution but it really does not matter, because different solutions give representations of the groups $\Gamma = \langle a,b,c\rangle$ which are conjugate in $\mathrm{PSL}(2,\mathbb{C})$. The last statement is a direct consequence of Lemma 3.1 since by the construction the different solutions correspond to the same $par(\Gamma)$. For definiteness we can always fix the analytic branches of the square roots in the following formulas.

Now we can write down the representation of $\langle a,b,c\rangle$ in $\mathrm{SL}(2,\mathbb{C})$:

$$A = \begin{pmatrix} 0 & i/\beta \\ i\beta & 0 \end{pmatrix},\quad B = \begin{pmatrix} 0 & i\beta \\ i/\beta & 0 \end{pmatrix},\quad C = \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & -c_{11} \end{pmatrix};$$

$$\beta = \sqrt{\frac{-\rho_0 + \sqrt{\rho_0^2 - 4}}{2}},$$

$$c_{21} = \frac{\rho_1/\beta - \rho_2\beta}{i/\beta^2 - i\beta^2},\quad c_{12} = \frac{-\rho_1\beta + \rho_2/\beta}{i/\beta^2 - i\beta^2},\quad c_{11} = i\sqrt{c_{12}c_{21}+1}; \tag{1}$$

$$\rho_k = -2\,cosh(\mu_k) \text{ for } k = 0,1,2.$$

To obtain the representation of $\langle a,b,c\rangle$ in $\mathrm{PSL}(2,\mathbb{C})$ one has to consider the image of the representation in $\mathrm{SL}(2,\mathbb{C})$ under canonical projection $p : \mathrm{SL}(2,\mathbb{C}) \to \mathrm{PSL}(2,\mathbb{C})$.

4. Deducing parameters from the singular structure

Knowing the structure of the singular set of a generalized triangle orbifold one can obtain a presentation of its fundamental group and a representation of the group in $\mathrm{SL}(2,\mathbb{C})$. This representation is defined by complex parameters ρ_0, ρ_1 and ρ_2, so it is possible to find the parameters, which correspond to a given singular structure. In this section and in Section 6 we show how it can be carried on for the common generalized triangle groups.

Suppose we are given a singular set of a common generalized triangle orbifold (Figure 3). Using Wirtinger algorithm we can start from the given three arcs a, b, c and passing through the tangles obtain words on a, b, c corresponding to all the other arcs. It is obvious that until we come to the central part of the diagram (D_1), we do not obtain any equations on the generators, so *all the words in the presentation of the group come from* D_1 (Figures 3, 5). Equations defining parameters are then obtained by taking traces of corresponding words in $\mathrm{SL}(2,\mathbb{C})$. With the help of the well known formulas

$$
\begin{aligned}
&\mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = \mathrm{tr}(\mathrm{W}_2\mathrm{W}_1),\\
&\mathrm{tr}(\mathrm{W}_1\mathrm{W}_2\mathrm{W}_1^{-1}) = \mathrm{tr}(\mathrm{W}_2),\\
&\mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = \mathrm{tr}(\mathrm{W}_1)\mathrm{tr}(\mathrm{W}_2) - \mathrm{tr}(\mathrm{W}_1^{-1}\mathrm{W}_2),\\
&\mathrm{tr}(\mathrm{H}^{-1}) = -\mathrm{tr}(\mathrm{H}) \quad \text{if H is a matrix of a half} - \text{turn}
\end{aligned}
\tag{2}
$$

the traces of the words on A, B, C can be expressed in terms of $\mathrm{tr(AB)} = \rho_0$, $\mathrm{tr(AC)} = \rho_1$, $\mathrm{tr(BC)} = \rho_2$ and $\mathrm{tr(ABC)}$ (an explicit expression for $\mathrm{tr(ABC)}$ in terms of ρ_i can be obtained using (1)). Since we have three complex parameters we always need three independent equations to define them. In the remaining part of this section we will give the presentations of the groups and corresponding equations for each of the cases $A - H$ from Theorem 2.4.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ w_1w_2,\ w_3w_4\rangle$,
equations:

$$
\left\{
\begin{array}{l}
\mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = 2,\\
\mathrm{tr}(\mathrm{W}_3\mathrm{W}_4) = 2,\\
\mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = 2
\end{array}
\right.
$$

(We denote by w_i words on a, b, c and by W_i — corresponding matrix words in $\mathrm{SL}(2,\mathbb{C})$, $i = 1, \ldots, 6$.)

It follows from [10] that one of the words w_1w_2, w_3w_4, w_5w_6 in the fundamental group is always a corollary of two other. However by the rigidity argument in the matrix group all the three words are independent and give three independent equations on the parameters. See Example 6A for the illustration.

B.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ w_1w_2,\ (w_3w_4)^t,\ (w_5w_6)^t \rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = 2, \\ \mathrm{tr}(\mathrm{W}_3\mathrm{W}_4) = -2\cos(\pi/\mathrm{t}), \\ \mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = -2\cos(\pi/\mathrm{t}). \end{cases}$$

C.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ (w_1w_2)^{t_1},\ (w_3w_4)^{t_2},\ (w_5w_6)^{t_3} \rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = -2\cos(\pi/\mathrm{t}_1), \\ \mathrm{tr}(\mathrm{W}_3\mathrm{W}_4) = -2\cos(\pi/\mathrm{t}_2), \\ \mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = -2\cos(\pi/\mathrm{t}_3). \end{cases}$$

D.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ (w_1w_2)^{t_1},\ (w_5w_6)^{t_2},\ (w_1w_2w_3)^{t_3} \rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = -2\cos(\pi/\mathrm{t}_1), \\ \mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = -2\cos(\pi/\mathrm{t}_3), \\ \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2\mathrm{W}_3) = -2\cos(\pi/\mathrm{t}_2). \end{cases}$$

It can be seen that word $W_1W_2W_3$ has an odd length, which follows that in this case we need the expression for tr(ABC). Actually, this is the only case where we may need the expression.

E.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ (w_1w_2)^2,\ (w_5w_6)^2 \rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = 0, \\ \mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = 0, \\ \mathrm{tr}(\mathrm{W}_3\mathrm{W}_4) = -2. \end{cases}$$

The third equation follows from the fact that the multiple of two half-turns with the parallel axes is a parabolic isometry, so w_3w_4 is a parabolic corresponding to the cusp. We choose sign $'-'$ for the trace of the parabolic element because the axes of the half-turns have the same directions.

F.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ w_1w_2\rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = 2, \\ \mathrm{tr}(\mathrm{W}_3\mathrm{W}_4) = -2, \\ \mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = -2. \end{cases}$$

G.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ (w_3w_4w_5)^2\rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_3\mathrm{W}_4\mathrm{W}_5) = 0, \\ \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = -2, \\ \mathrm{tr}(\mathrm{W}_3\mathrm{W}_4) = -2. \end{cases}$$

Here, as in case D, we again have a word $W_3W_4W_5$ of an odd length, but since $\mathrm{tr}(\mathrm{W}_3\mathrm{W}_4\mathrm{W}_5) = 0$ we can easily avoid using the explicit expression for tr(ABC) in this case. Really, using formulae (2) we can present $\mathrm{tr}(\mathrm{W}_3\mathrm{W}_4\mathrm{W}_5)$ as $\mathrm{tr(ABC)P[tr(AB)}$, $\mathrm{tr(AC), tr(BC)]}$ where P[...] is a polynomial, and since $\mathrm{tr(ABC)} \neq 0$ we obtain equation $\mathrm{P[tr(AB), tr(AC), tr(BC)]} = 0$. This procedure can be used for case H as well.

H.



Group:
$\langle a,\ b,\ c \mid a^2,\ b^2,\ c^2,\ (w_1w_2)^t,\ (w_1w_2w_3)^2\rangle$,
equations:

$$\begin{cases} \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2) = -2\cos(\pi/\mathrm{t}), \\ \mathrm{tr}(\mathrm{W}_5\mathrm{W}_6) = -2, \\ \mathrm{tr}(\mathrm{W}_1\mathrm{W}_2\mathrm{W}_3) = 0. \end{cases}$$

5. Connection with the two-generator Kleinian groups

Let $\Gamma_0 = \langle f, g\rangle$ be a discrete subgroup of $\mathrm{PSL}(2,\mathbb{C})$. There is a natural way to associate to Γ_0 a generalized triangle group Γ which is equal to Γ_0 or contains it as a subgroup of index 2. It follows that such properties as discreteness, arithmeticity and others of two-generator subgroups of $\mathrm{PSL}(2,\mathbb{C})$ can be studied using the generalized triangle groups.

The group Γ is constructed as follows (see [11]): If neither f nor g are parabolic then let N be a common perpendicular to the axes of f and g. In case of parabolic generator corresponding end of the line N is the fixed point of the parabolic. Obviously, this construction uniquely defines line N for any Kleinian group $\Gamma_0 = \langle f, g\rangle$. Let c be the half-turn about N. Then there exist such half-turns a and b that

$f = ac$ and $g = cb$ ([9], p. 47). So we obtain group $\Gamma = \langle a, b, c \rangle$ which contains Γ_0 as a subgroup of index at most 2.

Visa versa starting from the group Γ generated by three half-turns a, b, and c one can easily obtain its two generator subgroup $\Gamma_0 = \langle ab, ac \rangle$. Since $\Gamma = \Gamma_0 \cup a\Gamma_0$ ($a^2 = id$), index of Γ_0 in Γ is equal to 2 or 1 depending on if a is contained in Γ_0 or not.

The set of two-generator Kleinian groups is usually parameterized by three complex numbers [12]:

$$\beta(f) = \mathrm{tr}^2(\mathrm{f}) - 4,\ \beta(\mathrm{g}) = \mathrm{tr}^2(\mathrm{g}) - 4,\ \gamma(\mathrm{f}, \mathrm{g}) = \mathrm{tr}([\mathrm{f}, \mathrm{g}]) - 2.$$

A common computation with traces shows that this parameters are connected with our parameterization for the corresponding generalized triangle group by the following formulas:

$$\beta(f) = \rho_1^2 - 4,\ \beta(g) = \rho_2^2 - 4,\ \gamma(f, g) = \rho_0^2 + \rho_1^2 + \rho_2^2 + \rho_0\rho_1\rho_2 - 4.$$

Since we have described all the singular sets of the generalized triangle orbifolds we can at least theoretically present an algorithm, which using equations from the previous section enumerates all the parameters corresponding to the generalized triangle groups and discrete two-generator subgroups of $\mathrm{PSL}(2, \mathbb{C})$. The main practical difficulty is that the complexity of the equations grows as the singular structures become more complicated. The other point is that we have to cast away the redundant solutions of the equations somehow. In the next section we will show how this can be done for the examples considered, however we do not know any general method to carry on this procedure.

Let us also note that the suggested algorithm shows only that our set of parameters is semi-recursive: we can enumerate all the parameters of discrete generalized triangle groups, but there is no way to decide in a finite number of steps does a given triple of complex numbers define a discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$ or not.

6. EXAMPLES OF THE COMMON GENERALIZED TRIANGLE ORBIFOLDS.

Let us consider examples for different types of singularities from Theorem 2.4 and find parameters defining corresponding generalized triangle groups. Here we will try to give examples, which deserve a particular interest and have previously appeared in different situations. Detailed descriptions of some other generalized triangle orbifolds can be found in [2] and [3].

A.



$$a_{-1} = cac^{-1}$$
$$b_{-1} = aba^{-1}$$
$$c_{-1} = bcb^{-1}$$
$$a_1 = b_{-1}^{-1}ab_{-1}$$
$$b_1 = c_{-1}^{-1}bc_{-1}$$
$$b_2 = b_{-1}^{-1}b_1b_{-1}$$
$$a_{-1}^{-1}b_2 = ca^{-1}c^{-1}ab^{-1}a^{-1}bc^{-1}bcb^{-1}aba^{-1}$$

Matrix group:

$$\begin{gathered}\langle A,\ B,\ C \mid -A^2, -B^2, -C^2,\\ CA^{-1}C^{-1}AB^{-1}A^{-1}BC^{-1}BCB^{-1}ABA^{-1},\\ AB^{-1}A^{-1}BC^{-1}B^{-1}CA^{-1}CAC^{-1}BCA^{-1},\\ BC^{-1}B^{-1}CA^{-1}C^{-1}AB^{-1}ABA^{-1}CAB^{-1}\rangle\end{gathered}$$

Let tr(AB) = t_0, tr(AC) = t_1, tr(BC) = t_2.
By taking traces of the words we obtain a system of equations on t_0, t_1 and t_2[1]:

$$\begin{cases} 3t_0 - t_0^3 - 2t_0t_1^2 + t_0^3t_1^2 + 2t_0^2t_1t_2 - t_0^2t_1^3t_2 - 2t_0t_2^2 + t_0^3t_2^2 - t_0^3t_1^2t_2^2 - t_0^2t_1t_2^3 = 2,\\ 3t_2 - t_2^3 - 2t_2t_0^2 + t_2^3t_0^2 + 2t_2^2t_0t_1 - t_2^2t_0^3t_1 - 2t_2t_1^2 + t_2^3t_1^2 - t_2^3t_0^2t_1^2 - t_2^2t_0t_1^3 = 2,\\ 3t_1 - t_1^3 - 2t_1t_2^2 + t_1^3t_2^2 + 2t_1^2t_2t_0 - t_1^2t_2^3t_0 - 2t_1t_0^2 + t_1^3t_0^2 - t_1^3t_2^2t_0^2 - t_1^2t_2t_0^3 = 2. \end{cases}$$

We are interested only in complex solutions of this system and by the symmetry of the singular graph all the three roots should be equal. The only such solutions are (approximately):

$$(t_0, t_1, t_2) = (0.662359 \pm 0.56228i, 0.662359 \pm 0.56228i, 0.662359 \pm 0.56228i).$$

So we have $\rho_0 = \rho_1 = \rho_2 = 0.662359\dots \pm 0.56228\dots i$. The signs before the imaginary part correspond to the orientations of the generating arcs and do not affect the presentation of the group in $\mathrm{PSL}(2,\mathbb{C})$.

The primitive polynomial of the root is $t^3 - t + 1$, using the arithmeticity test for generalized triangle groups from [3] it can be easily verified that the orbifold is arithmetic. It is also interesting to note (see [26]) that the two-fold covering of this orbifold is Weeks-Mateveev-Fomenko manifold — the hyperbolic manifold of the smallest known volume.

(Of course, here we could use the symmetry of the singular graph from the beginning and initially suppose tr(AB) = tr(AC) = tr(BC). It would considerably simplify the calculations but leads to some loss of generality.)

---

[1] We use an original computer program <tracer>(C++, 242 lines) to calculate trace polynomials for generalized triangle groups. The program is available on request.



$$\begin{aligned} a_1 &= bab^{-1} \\ b_1 &= cbc^{-1} \\ c_1 &= aca^{-1} \\ a_{-1} &= c_1^{-1}ac_1 \\ b_{-1} &= a_1^{-1}ba_1 \end{aligned}$$

$$\begin{aligned} a_{-1}^{-1}a_1 &= ac^{-1}a^{-1}ca^{-1}bab^{-1} \\ b_{-1}^{-1}c^{-1} &= ba^{-1}b^{-1}ab^{-1}c^{-1} \\ b_1c_1 &= cbc^{-1}aca^{-1} \end{aligned}$$

Equations on $t_0 = \mathrm{tr(AB)}$, $t_1 = \mathrm{tr(AC)}$, $t_2 = \mathrm{tr(BC)}$:

$$\begin{cases} t_0^2t_1^2 + t_0t_1t_2 - t_0^2 - t_1^2 + 2 = 2, \\ t_0^2t_2 + t_0t_1 - t_2 = -2cos(\pi/n), \\ t_1^2t_2 + t_0t_1 - t_2 = -2cos(\pi/n). \end{cases}$$

We will consider only $n = 3$. Up to the symmetry and change of orientation the system has only one complex solution, which defines the complex place of the orbifold group:

$$\rho_0 = \frac{i+\sqrt{3}}{2},\ \rho_1 = \frac{i-\sqrt{3}}{2},\ \rho_2 = 0.$$

The squares of ρ_0, ρ_1 are algebraic integers and by arithmeticity test [3] the group is arithmetic defined over field $\mathbb{Q}[\sqrt{-3}]$.
(For $n = 2$ and $n = \infty$ the group has no any complex places, so the orbifold is not hyperbolic. For large values of n the group has more then one complex places induced by $cos(\pi/n) \to cos(k\pi/n)$, $(k, n) = 1$ and so it is not arithmetic.)

C.



$$b_{-2} = b^{-1}aba^{-1}b$$
$$b_{-1} = aba^{-1}$$
$$b_1 = b_{-2}bb_{-2}^{-1}$$
$$b_2 = cb_1c^{-1}$$
$$a_1 = b_{-1}^{-1}ab_{-1}$$
$$c_1 = aca^{-1}$$

$$c_1a_1 = acb^{-1}aba^{-1}$$
$$b_{-2}^{-1}c^{-1} = b^{-1}ab^{-1}a^{-1}bc^{-1}$$
$$b_2a^{-1} = cb^{-1}aba^{-1}bab^{-1}a^{-1}bc^{-1}a^{-1}$$

Equations on $t_0 = \mathrm{tr(AB)}$, $t_1 = \mathrm{tr(AC)}$, $t_2 = \mathrm{tr(BC)}$:

$$\begin{cases} t_0t_2 + t_1 = 0, \\ t_0^2t_2 + t_0t_1 - t_2 = -2cos(\pi/n), \\ (t_0^3 - 3t_0)(t_0t_1t_2 + t_1^2 + t_0^2 - 2) + t_0t_1^2 + t_1t_2 - t_0 = 0. \end{cases}$$

Complex solution ($n > 3$):

$$\rho_0^2 = \tfrac{1}{2}(1 + 4sin^2(\pi/n)(1 \pm \sqrt{1 - \tfrac{4}{1+4sin^2(\pi/n)}}),$$
$$\rho_1 = -2cos(\pi/n)\rho_0,$$
$$\rho_2 = 2cos(\pi/n).$$

(Here ρ_2 is always a real number; for $n = 2, 3$ ρ_0 and ρ_1 are also real, so corresponding orbifolds are not hyperbolic 3-orbifolds of finite volume. According to [8] for $n = 2$ the orbifold admits the spherical structure and for $n = 3$ it is Euclidean.)

As it was shown in [27] the considered orbifolds can be obtained as the factor-orbifolds of Fibonacci manifolds $F(2n, n)$ by their full groups of isometries. So using the arithmeticity test from [3] we can now easily find all arithmetic Fibonacci manifolds (see also [20]). The field of definition of the orbifold group $\mathbb{Q}[\rho_0^2, cos^2(\pi/n)]$ has exactly one complex place only if $n =$ 4, 5, 6, 8 12, ∞; it is a matter of direct verification that this cases satisfy the remaining conditions of the arithmeticity test. Thus the orbifolds and corresponding Fibonacci manifolds are arithmetic for $n =$ 4, 5, 6, 8 12, ∞.



$$\begin{aligned}
c_{-2} &= ac_{-1}a^{-1}\\
c_{-1} &= b^{-1}cb\\
b_{-1} &= c_{-2}^{-1}bc_{-2}\\
a_1 &= c_{-2}^{-1}ac_{-2}\\
b_1 &= cbc^{-1}\\
c_1 &= b_1cb_1^{-1}
\end{aligned}$$

$$\begin{aligned}
a_1b_{-1}^{-1} &= ab^{-1}c^{-1}ab^{-1}cba^{-1}\\
c_{-2}^{-1}c_1 &= ab^{-1}c^{-1}ba^{-1}cbcb^{-1}c^{-1}\\
a^{-1}a_1b_{-1}^{-1} &= b^{-1}c^{-1}ab^{-1}cba^{-1}
\end{aligned}$$

Equations on $t_0 = \mathrm{tr}(\mathrm{AB})$, $t_1 = \mathrm{tr}(\mathrm{AC})$, $t_2 = \mathrm{tr}(\mathrm{BC})$:

$$\begin{cases} t_0 = 0,\\ 4t_2^2 - t_1^2t_2^2 - t_2^4 + t_1^2 - 2 = 0,\\ 4t_1^2 - t_1^2t_2^2 - t_1^4 = 4cos^2(\pi/n). \end{cases}$$

Equation $\mathrm{tr}(\mathrm{AB}) = 0$ geometrically means that the axes of half-turns a and b intersect and orthogonal in $\mathcal{H}^3$.

The system has exactly one pair of complex-conjugate roots for any $n > 2$ (for $n = 2$ the orbifold is Euclidean), this roots give the values of parameters $\rho_0(= 0)$, ρ_1, ρ_2 defining the orbifold group. Since the analytic formulas for the solutions are rather complicated we do not give them here. One can see that the field of definition $\mathbb{Q}[\rho_1^2, \rho_2^2]$ of the group has exactly one complex place iff $n = 3$, 4, 6, and it is easy to check the other conditions of the arithmeticity test [3] for this values of n. So we obtain, that the orbifolds are arithmetic for $n = 3$, 4, 6.

This groups and orbifolds first appeared in [18] as an infinite one-parameter family extending one of the three regular tessellations of $\mathcal{H}^3$. It is interesting to note that the corresponding family for the one of the two remaining regular tessellations is also known [17] and consists of Fibonacci manifolds considered in the previous example; and, finally, for the third tessellation the question of constructing such a family is still open.

E.



$$\begin{aligned} a_1 &= bab^{-1} \\ b_1 &= a_1 b a_1^{-1} \\ c_1 &= aca^{-1} \\ a_2 &= c a_1 c^{-1} \end{aligned}$$

$$\begin{aligned} c_1 b^{-1} &= aca^{-1}b^{-1} \\ a_2 a^{-1} &= cbab^{-1}c^{-1}a^{-1} \\ b_1 c^{-1} &= baba^{-1}b^{-1}c^{-1} \end{aligned}$$

Equations on $t_0 = \mathrm{tr(AB)}$, $t_1 = \mathrm{tr(AC)}$, $t_2 = \mathrm{tr(BC)}$:

$$\begin{cases} t_0 t_1 + t_2 = 0, \\ t_0 t_1 t_2 + t_0^2 + t_1^2 - 2 = 0, \\ -t_0^2 t_2 - t_0 t_1 + t_2 = -2. \end{cases}$$

Solution:

$$\rho_0^2 = \rho_1^2 = 1 + i, \ \rho_2^2 = 2i.$$

This orbifold first appeared in [1] as the third hyperbolic 3-orbifold with a non-rigid cusp. We see that the orbifold is arithmetic with the field of definition $\mathbb{Q}[i]$ and so its group is commensurable in $\mathrm{PSL}(2, \mathbb{C})$ with Picard group $\mathrm{PSL}(2, \mathbb{Z}[\mathrm{i}])$.

F.



$a_1 = bab^{-1}$
$a_2 = ca_1c^{-1}$
$b_1 = cbc^{-1}$
$b_2 = a_2b_1a_2^{-1}$
$b_3 = b^{-1}b_2b$
$c_{-1} = b^{-1}cb$
$c_{-2} = b_3^{-1}c_{-1}b_3$
$b_{-1} = a_2^{-1}ba_2$

$c_{-2}^{-1}a_2 =$
$b^{-1}cbab^{-1}a^{-1}b^{-1}c^{-1}baba^{-1}b^{-1}c^{-1}bcbab^{-1}c^{-1}$
$b_{-1}^{-1}c = cba^{-1}b^{-1}c^{-1}b^{-1}cbab^{-1}$
$a^{-1}b_3 = a^{-1}b^{-1}cbaba^{-1}b^{-1}c^{-1}b$

Equations on $t_0 = \mathrm{tr(AB)}$, $t_1 = \mathrm{tr(AC)}$, $t_2 = \mathrm{tr(BC)}$:

$$\begin{cases} t_0^2t_1^3t_2^2 + t_0t_1^2t_2(2 - 2t_2^2 + t_0^2(-2 + 3t_2^2)) + t_0t_2(5 - 4t_2^2 + t_2^4 + t_0^4(-1 + t_2^2)^2 + \\ \quad +t_0^2(-4 + 5t_2^2 - 2t_2^4)) + t_1(1 - 3t_2^2 + t_2^4 + t_0^2(-3 + 7t_2^2 - 4t_2^4) + t_0^4(1 - 4t_2^2 + 3t_2^4)) = 2, \\ t_2 + (t_1 + t_0t_2)(t_0 - t_1t_2 - t_0t_2^2) = -2, \\ t_0 + (t_1 + t_2t_0)(t_2 - t_1t_0 - t_2t_0^2) = -2. \end{cases}$$

Complex solution:

$$\rho_1 = 2,\ \rho_0 = \rho_2 = \frac{-1 \pm \sqrt{-3}}{2}.$$

This example appeared as an orbifold that has the sister-figure-eight knot as a two-shit cover. As we see, the orbifold is arithmetic (by test [3]) with the field of definition $\mathbb{Q}[\sqrt{-3}]$.

G.



$a_1 = bab^{-1}$
$b_1 = cbc^{-1}$
$c_1 = aca^{-1}$
$a_{-1} = c_1^{-1}ac_1$
$c_{-1} = a_1ca_1^{-1}$

$a_1b_1c_1 = bab^{-1}cbc^{-1}aca^{-1}$
$b^{-1}c_{-1}^{-1} = ab^{-1}c^{-1}ba^{-1}b^{-1}$
$a_1b_1 = bab^{-1}cbc^{-1}$

With our usual notation $t_0 = \mathrm{tr(AB)}$, $t_1 = \mathrm{tr(AC)}$, $t_2 = \mathrm{tr(BC)}$ we obtain $\mathrm{tr(A_1B_1C_1)} = \mathrm{tr(ABC)(t_0t_1t_2 + 1)} = 0$. Since $\mathrm{tr(ABC)} \neq 0$ it gives $t_0t_1t_2 + 1 = 0$, and so we have the following system of equations on the parameters:

$$\begin{cases} t_0t_1t_2 = -1, \\ t_0t_2^2 + t_1t_2 - t_0 = -2, \\ t_0^2t_2 + t_1t_0 - t_2 = -2. \end{cases}$$

Complex solution:

$$\rho_0 = \rho_1 = \rho_2 = \frac{1 \pm \sqrt{-3}}{2}.$$

Since $\rho_0 = \rho_1 = \rho_2$ the orbifold has an order 3 symmetry, which cyclically changes the generators of the group. This symmetry can be easily seen on the spacial representation of the singular graph but is lost on the plane projection. According to [3] the considered orbifold is one of the three arithmetic non-compact generalized triangle orbifolds with equal parameters (the remaining two orbifolds correspond to Case C).

H. We have tried more then 20 different structures for this case but did not succeed in finding any arithmetic examples. The central part structure implies rather strong restrictions on the orbifold: it should be non compact and it is essentially asymmetric. Let us leave as a challenge *to find an arithmetic example for case H or prove that it does not exist.*

7. An example of the exceptional generalized triangle orbifold.

Here we will consider an example of a generalized triangle orbifold, whose singular set does not correspond to any of the eight types of singularities from Theorem 2.4. This orbifold appears to be a counterexample to the Lemma 2.3 if we do not suppose that the orbifolds considered there are common.

The example is well known, it is the Picard orbifold. It was first shown in [6] that the Picard group can be generated by only two elements or by three involutions. Let us reproduce what happens.



$b_1 = cbc^{-1}$
$b_{-1} = aba^{-1}$

We won't find the matrix representation in $\mathrm{SL}(2, \mathbb{C})$ for the Picard group since it is well known and can be easily obtained from the above. Our goal is only to show that this group can be generated by three involutions, so we will make all the calculations in the group, which is a bit easier then working with the matrices.

We have

$$\Gamma = < a, b, c, x \mid a^2, b^2, c^2, x^2, (xac)^2, (xabc)^3, (bax)^3, (abax)^2 > .$$

$(xac)^2 = 1$ implies $acx = xca$;
$(abax)^2 = 1$implies $xaba = abax$.

Now consider the word $xabc$:

$$1 = (xabc)^3 = xabcxabcxabc = xabcxab\ aa\ cxabc = xabc\ abax\ xca\ abc = xabcabacbc;$$

$$x = abcabacbc.$$

So $\Gamma = < a, b, c \mid a^2, b^2, c^2, (abcabacbc)^2, (abcabacbcac)^2, (abacb)^3, (acabacbc)^2 > .$

Note, that in the notations of Lemma 2.3 here we have $H_1(\tilde{X}) \cong \mathbb{Z}^4$ (so the rank of $\pi_1(\tilde{X})$ is greater then 3 and the Picard group can not be a common generalized triangle group by the definition) but, the same time, $H_1(X) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ has rank 2.

The considered example is non compact. One can obtain an infinite series of the compact exceptional generalized triangle orbifolds by $(n; 0)$ Dehn surgeries on the cusp of the Picard orbifold.

References

[1] C. Adams, Noncompact hyperbolic 3-orbifolds of small volume, In: *Topology'90*, edited by B. Apanasov, W. D. Neumann, A. W. Reid, and L. Siebenmann, de Gruyter, Berlin (1992), 1–15.

[2] M. Belolipetsky, Arithmetic properties of the groups of isometries generated by three involutions (in Russian), *Magister Thesis*, Novosibirsk State University, Novosibirsk (1997), 30 pp.

[3] M. Belolipetsky, On arithmetic Kleinian groups generated by three half-turns, Preprint 00-010, Universität Bielefeld, Bielefeld (2000), 17 pp.

[4] A. Berdon, The geometry of discrete groups, Springer-Verlag, 1983.

[5] R. Brooks, J. P. Matelski, The dynamics of 2-generator subgroups of PSL(2, ℂ), *Riemann surfaces and related topics: Proc. of the 1978 Stony Brook Conf.* (I. Kra and B. Maskit, eds), 65–71.

[6] A. M. Brunner, A two-generator presentation for the Picard group. *Proc. of Amer. Math. Soc.*, 115, N 1 (1992), 45–46.

[7] M. D. E. Condor, C. Maclachlan, G. Martin, E. A. O'Brien, 2-generator arithmetic Kleinian groups III, *DVI preprint*, Department of Mathematical Sciences, University of Aberdeen.

[8] W. Dunbar, Geometric orbifolds, *Rev. Mat. Univ. Complutense Madr.*, 1 (1988), 67–99.

[9] W. Fenchel, Elementary geometry in hyperbolic space, De Gruyter, 1989.

[10] R. Fox, A quick trip through knot theory, Topology of 3-manifolds, *Proc. 1961 Top. Inst. Univ. Georgia*, Prentice Hall.

[11] J. Gilman, A discreteness condition for subgroups of PSL(2, ℂ), *Contemp. Math.*, 211 (1997), 261–267.

[12] F. Gehring and G. Martin, Commutators, collars and the geometry of Möbius groups, *J. d'Analyse Math.*, 63 (1994), 175–219.

[13] F. Gehring, J. Gilman, G. Martin, Discrete groups with real parameters, *preprint*, 1998.

[14] F. Gehring, C. Maclachlan, G. Martin, Two-generator arithmetic Kleinian groups II, *Bull. Lond. Math. Soc.*, 30, N 3 (1998), 258–266.

[15] F. Gehring, C. Maclachlan, G. Martin, A. Reid, Arithmeticity, Discretness and Volume, *Trans. Am. Math. Soc.*, 349, N 9 (1997), 3611–3643.

[16] A. Haefliger, N. D. Quach, Une presentation de groupe fundamental d'une orbifold, *Asterisque*, 116 (1984), 98–107.

[17] H. Helling, A. C. Kim, J. L. Mennicke, A geometric study of Fibonacci groups, *J. Lie Theory*, 8, N 1 (1998), 1–23.

[18] H. Helling, A. C. Kim, J. L. Mennicke, Some honey-combs in hyperbolic 3-space, *Comm. in Algebra*, 23 (1995), 5169–5206.

[19] H. M. Hilden, M.-T. Lozano and J. M. Montesinos, A characterisation of arithmetic subgroups of $\mathrm{SL}(2,\mathbb{R})$ and $\mathrm{SL}(2,\mathbb{C})$, *Math. Nachr.*, 159 (1992), 245–270.

[20] H. M. Hilden, M.-T. Lozano, J. M. Montesinos, The arithmeticity of the figure-eight knot orbifolds, In: *Topology'90*, edited by B. Apanasov, W. D. Neumann, A. W. Reid, and L. Siebenmann, de Gruyter, Berlin (1992), 169–183.

[21] T. Jørgensen, On discrete groups of Möbius transformations. *Amer. J. Math.*, 78 (1976), 739–749.

[22] E. Klimenko, Discrete groups in three-dimensional Lobachevsky space generated by two rotations, *Sib. Math. J.*, 30 (1989), 95–100.

[23] E. Klimenko, Some remarks on subgroups of $\mathrm{PSL}(2,\mathbb{C})$, *Q & A in General Topology*, 8, N 2 (1990), 371–381.

[24] E. Klimenko, N. Kopteva, Discreteness criteria for $\mathcal{RP}$-groups, *Preprint 00-061*, Universität Bielefeld, Bielefeld (2000), 22 pp.

[25] C. Maclachlan, G. Martin, 2-generator arithmetic Kleinian groups, *J. Reine Angew. Math.*, 511 (1999), 95–117.

[26] A. Mednykh, A. Vesnin, Visualization of the isometry group action on the Fomenko-Matveev-Weeks manifold, *J. Lie Theory*, 8, N 1 (1998), 51–66.

[27] A. Rasskazov, A. Vesnin, Isometries of hyperbolic Fibonacci manifolds, *Sib. Math. J.*, 40 (1999), 9–22.

[28] W. Thurston, The geometry and topology of 3-manifolds, Lecture Notes, Princeton University, 1980.

Sobolev Institute of Mathematics, Novosibirsk, 630090, Russia
e-mail: `mbel@math.nsc.ru`